

Determination of α_s from Event Shapes and Power Corrections

Daniel Wicke^a

^aFachbereich Physik, Bergische Universität GH, Gaußstr. 20, 42097 Wuppertal; wicke@cern.ch

The size of non-perturbative corrections to event shape observables is predicted to fall like a power of the inverse centre of mass energy. These power corrections are investigated for different observables from e^+e^- -annihilation and compared to the theoretical predictions. Using the latest DELPHI high energy data advantages of determining α_s from these predictions are discussed and compared to conventional methods.

1. Introduction

The upgrade of LEP to run at energies above the Z -peak induced new possibilities in measuring the running of α_s . Due to asymptotic freedom, non-perturbative effects become less important at higher energies. Different theoretical frameworks suggest that for event shapes the non-perturbative corrections reduce like the inverse power of the centre of mass energy [1,2]. Theoretical predictions for this decrease can be used in the measurement of α_s independent of event by event hadronization models.

2. α_s from Mean Event Shapes using Power Corrections

The analytical power ansatz for non-perturbative corrections by Dokshitzer and Webber [2,3] can be used to determine α_s from mean event shapes [4–6]. This ansatz provides an additive term to the perturbative $\mathcal{O}(\alpha_s^2)$ QCD prediction $\langle f_{\text{pert}} \rangle$:

$$\langle f \rangle = \frac{1}{\sigma_{\text{tot}}} \int f \frac{df}{d\sigma} d\sigma = \langle f_{\text{pert}} \rangle + \langle f_{\text{pow}} \rangle \quad (1)$$

The power correction is given by

$$\langle f_{\text{pow}} \rangle = a_f \cdot \frac{\mu_I}{E_{\text{cm}}} \left[\bar{\alpha}_0(\mu_I) - \alpha_s(\mu) - \left(\beta_0 \cdot \ln \frac{\mu^2}{\mu_I^2} + \frac{K}{2\pi} + 2\beta_0 \right) \alpha_s^2(\mu) \right] \quad (2)$$

$\bar{\alpha}_0$ is a non-perturbative parameter accounting for the contributions to the event shape below

an infrared matching scale μ_I , $K = (67/18 - \pi^2/6)C_A - 5N_f/9$, $\beta_0 = (33 - 2N_f)/12\pi$ and $a_f = 4C_f/\pi$. Beside α_s this formulae contains $\bar{\alpha}_0$ as the only free parameter. In order to measure α_s this parameter has to be known.

To infer $\bar{\alpha}_0$ a combined fit of α_s and $\bar{\alpha}_0$ to a large set of measurements at different energies [4,5,7] is performed. For $E_{\text{cm}} \geq M_Z$ only DELPHI measurements are included in the fit. Fig. 2 shows the measured mean values of $\langle 1 - T \rangle$ and $\langle M_h^2/E_{\text{vis}}^2 \rangle$ as a function of the centre of mass energy together with the results of the fit. The resulting values of $\bar{\alpha}_0$ are summarized in Tab. 1. The extracted $\bar{\alpha}_0$ values are around 0.5 as expected in [3], but they are incompatible with each other. As the assumed universality is not given to the precision that is accessible from data, $\bar{\alpha}_0$ is individually determined for $\langle 1 - T \rangle$ and $\langle M_h^2/E_{\text{vis}}^2 \rangle$. The scale error is obtained from varying the renormalization scale $f = \mu^2/E_{\text{cm}}^2$ from 0.25 to 4 and the infrared matching scale from 1GeV to 2GeV.

After fixing $\bar{\alpha}_0$, the α_s values corresponding to the high energy data points can be calculated from Eqs. (1–2). α_s is calculated for both observables individually and then combined with an

Table 1

$\bar{\alpha}_0$ as determined from mean event shapes.

Observable	$\bar{\alpha}_0 \pm \text{stat.} \pm \text{scale}$	χ^2/ndf
$\langle 1 - T \rangle$	$0.531 \pm 0.012 \pm 0.003$	42/23
$\langle M_h^2/E_{\text{vis}}^2 \rangle$	$0.434 \pm 0.010 \pm 0.010$	4.0/14

Table 2

α_s as obtained from means with the Eqs. (1-2) by averaging the $\langle 1 - T \rangle$ and $\langle M_h^2/E_{\text{vis}}^2 \rangle$ results.

E_{cm}	$\alpha_s(E_{\text{cm}}) \pm \text{stat.} \pm \text{sys.} \pm \text{scale}$
133GeV	0.1162 $\pm 0.0070 \pm 0.0009 \pm 0.0047$
161GeV	0.1047 $\pm 0.0069 \pm 0.0034 \pm 0.0043$
172GeV	0.1052 $\pm 0.0082 \pm 0.0037 \pm 0.0041$

unweighted average. Its error is propagated from the data and combined by assuming full correlation. An additional scale error was calculated by varying f and μ_I in the ranges discussed. The results are summarized in Tab. 2 and plotted as function of E_{cm} (open dots) together with the QCD expectation in Fig. 2a.

3. α_s from Event Shapes Distributions

From event shape distributions α_s is determined by fitting an α_s dependent QCD prediction ($\mathcal{O}(\alpha_s^2)$, pure NLLA or combined $\mathcal{O}(\alpha_s^2)$ +NLLA [8-10]) folded with a hadronization or a power correction to the data. The fit ranges used for the different QCD predictions are shown in Fig. 1. The ranges for pure NLLA and $\mathcal{O}(\alpha_s^2)$ fits are chosen to be distinct, so that the results are statistically uncorrelated.

3.1. Using Hadronization Models

As an example of a Monte Carlo based model in this section the hadronization correction is calculated using the JETSET PS model (Version 7.4 tuned to DELPHI data [11]). The QCD prediction is multiplied in each bin by the correction factor (the model prediction of the ratio of hadron over parton level in that bin).

For $\mathcal{O}(\alpha_s^2)$ -fits the renormalization scale μ was fitted to the LEP1 data [11] for both observables individually and then fixed for the fits to the high energy distributions. This takes advantage of optimized scales. In contrast for NLLA and combined fits μ was set equal to E_{cm} , so that these results can be compared to other experiments.

The systematic errors were obtained by fitting




Figure 1. Fit ranges chosen for fitting α_s from different QCD predictions of $1 - T$ and M_h^2/E_{vis}^2 distributions.

several $1 - T$ and M_h^2/E_{vis}^2 distributions evaluated by applying different cuts. The mean deviation from the central value is used as systematic error. Scale errors were taken from previous DELPHI publications [12,13].

The α_s values evaluated from the distributions are given in Tab. 3 and shown as a function of E_{cm} in Fig. 2b. The results agree well with those measured from the event shape means.

3.2. Using Power Corrections

Dokshitzer and Webber recently extended their model of power corrections to event shape distributions [14]. For distributions the power corrections are applied by shifting the complete prediction:

$$F(f) = \frac{1}{\sigma_{\text{tot}}} \frac{d\sigma}{df} = F_{\text{pert}}(f - \delta f) \quad (3)$$

where the size of the shift is given by the size of the power correction to the corresponding mean ($\delta f = \langle f_{\text{pow}} \rangle$). The perturbative prediction F_{pert} can now be any of the available QCD predictions: $\mathcal{O}(\alpha_s^2)$, NLLA or any combined scheme.

To measure α_s at a single energy, again one first has to determine $\bar{\alpha}_0$. This is done in a combined fit of $\bar{\alpha}_0$ and α_s to measurements [4,5,7] at many different energies. See Fig. 3a, c and Tab. 4. For M_h^2/E_{vis}^2 some of the lowest energies were re-




Figure 2. a) Measured mean values of $\langle 1 - T \rangle$ and $\langle M_h^2/E_{vis}^2 \rangle$ as a function of the centre of mass energy. The solid lines present the results of the fits with Eqs. (1–2), the dashed lines show the perturbative part. b) Energy dependence of α_s . The errors shown are statistical only. The band shows the QCD expectation of extrapolating the world average to other energies. c) α_s determined using Dokshitzer and Webber power corrections with $1 - T$ distribution.

moved from the fit, because the shifted QCD prediction was no longer defined at the lower end of the fit range.

To compare the quality of these fits to a hadronization model based description, a simultaneous fit of α_s to these data using the JETSET PS based hadronization correction is done. The results are plotted in Fig. 3b, d and summarized in Tab. 4.

Using these $\bar{\alpha}_0$ results to determine α_s from individual high energy $1 - T$ distributions lead to systematic deviations from the QCD expectation (See Fig. 2c). The deviation is different for NLLA and $\mathcal{O}(\alpha_s^2)$ (which use different fit ranges) and opposite in sign.

To investigate this problem, the Dokshitzer and Webber power correction term was replaced by a simple ansatz:

$$\delta f = \frac{C_1}{E_{cm}} + \frac{C_2}{E_{cm}^2} \quad (4)$$

A combined fit of C_1 , C_2 to the complete set of distributions probes the need of the quadratic power correction term. α_s was fixed to 0.120 here. It is found, that for the fit ranges used in $\mathcal{O}(\alpha_s^2)$ and NLLA fits of Thrust the correction arising from the quadratic power term is not negligible (See Tab. 5). The small result of C_2 for the combined result thus may be accidental.

4. Summary/Conclusions

Several methods of determining α_s at high energies from event shapes were discussed.

The results from means using power corrections are compatible with those from distributions using standard hadronization corrections. Results from power corrections yield smaller scale errors.

Power corrections to event shape distributions are more tricky.




Figure 3. $\mathcal{O}(\alpha_s^2)$ fit of α_s (and $\bar{\alpha}_0$) to $1-T$ and M_h^2/E_{vis}^2 distributions measured at different energies. In plot a,c the power ansatz Eq. 3, in b,d JETSET PS based correction is used to include non-perturbative contributions. The distributions including dotted lines were not included in the fit.

Table 3
 α_s as obtained from distributions using JETSET.

133GeV	$\alpha_s(E_{\text{cm}}) \pm$	exp.	$\pm$ scale
$\mathcal{O}(\alpha_s^2)$	$0.1154 \pm$	0.0072	± 0.006
NLLA	$0.1196 \pm$	0.0129	± 0.005
Combined	$0.1182 \pm$	0.0083	± 0.006
161GeV	$\alpha_s(E_{\text{cm}}) \pm$	stat. $\pm$ sys.	$\pm$ scale
$\mathcal{O}(\alpha_s^2)$	$0.1084 \pm 0.0088 \pm 0.0049 \pm 0.006$		
NLLA	$0.1056 \pm 0.0096 \pm 0.0043 \pm 0.005$		
Combined	$0.1120 \pm 0.0074 \pm 0.0032 \pm 0.006$		
172GeV	$\alpha_s(E_{\text{cm}}) \pm$	stat. $\pm$ sys.	$\pm$ scale
$\mathcal{O}(\alpha_s^2)$	$0.1151 \pm 0.0082 \pm 0.0058 \pm 0.005$		
NLLA	$0.0976 \pm 0.0097 \pm 0.0046 \pm 0.005$		
Combined	$0.1097 \pm 0.0076 \pm 0.0029 \pm 0.005$		

Table 4
 $\bar{\alpha}_0$ as determined from event shape distributions.
The error is the statistical error from the fit.

$1 - T$	$\bar{\alpha}_0$	χ^2/ndf
$\mathcal{O}(\alpha_s^2)$	0.550 ± 0.012	120/107
NLLA	0.475 ± 0.006	81/26
Combined	0.491 ± 0.005	377/137
M_h^2/E_{vis}^2	$\bar{\alpha}_0$	χ^2/ndf
$\mathcal{O}(\alpha_s^2)$	0.698 ± 0.014	83/49
NLLA	0.480 ± 0.012	3.2/7
Combined	0.554 ± 0.005	285/53

Table 5
Results of fitting the simple power ansatz to event shape distributions. $\alpha_s=0.120$.

$1 - T$	$C_1[\text{GeV}]$	$C_2[\text{GeV}^2]$
$\mathcal{O}(\alpha_s^2)$	1.23 ± 0.09	-10.4 ± 1.3
NLLA	0.27 ± 0.02	$+10.0 \pm 0.6$
Combined	0.73 ± 0.02	-0.48 ± 0.61
M_h^2/E_{vis}^2	$C_1[\text{GeV}]$	$C_2[\text{GeV}^2]$
$\mathcal{O}(\alpha_s^2)$	0.27 ± 0.09	$+14.8 \pm 6.9$
NLLA	not converging	
Combined	0.70 ± 0.02	-4.5 ± 6.9

REFERENCES

1. V. I. Zakharov. In these proceedings.
2. Y. L. Dokshitzer and B. R. Webber. *Phys. Lett.* **B352**(1995) 451.
3. B. R. Webber. Talk given at workshop on DIS and QCD in Paris, hep-ph/9510283, 1995.
4. DELPHI Coll., P. Abreu et al. *Z. Phys.* **C573**(1997) 229.
5. J. Drees, A. Grefrath, K. Hamacher, O. Passon, and D. Wicke. DELPHI 97-92 CONF 77, contrib. to EPS HEPC in Jerusalem, 1997.
6. O. Biebel. In these proceedings.
7. ALEPH Coll., *Phys. Lett.* **B284** (1992) 163. ALEPH Coll., *Z. Phys.* **C55** (1992) 209. AMY Coll., *Phys. Rev. Lett.* **62** (1989) 1713. AMY Coll., *Phys. Rev.* **D41** (1990) 2675. CELLO Coll., *Z. Phys.* **C44** (1989) 63. HRS Coll., *Phys. Rev.* **D31** (1985) 1. JADE Coll., *Z. Phys.* **C25** (1984) 231. JADE Coll., *Z. Phys.* **C33** (1986) 23. L3 Coll., *Z. Phys.* **C55** (1992) 39. Mark II Coll., *Phys. Rev.* **D37** (1988) 1. Mark II Coll., *Z. Phys.* **C43** (1989) 325. MARK J Coll., *Phys. Rev. Lett.* **43** (1979). OPAL Coll., *Z. Phys.* **C59** (1993) 1. PLUTO Coll., *Z. Phys.* **C12** (1982) 297. SLD Coll., *Phys. Rev.* **D51** (1995) 962. TASSO Coll., *Phys. Lett.* **B214** (1988) 293. TASSO Coll., *Z. Phys.* **C45** (1989) 11. TASSO Coll., *Z. Phys.* **C47** (1990) 187. TOPAZ Coll., *Phys. Lett.* **B227** (1989) 495. TOPAZ Coll., *Phys. Lett.* **B278** (1992) 506. TOPAZ Coll., *Phys. Lett.* **B313** (1993) 475.
8. R. K. Ellis, D. A. Ross, and A. E. Terrano. *Nucl. Phys.* **B178**(1981) 412.
9. S. Catani, G. Turnock, B. R. Webber, and L. Trentadue. *Phys. Lett.* **B263**(1991) 491.
10. S. Catani, G. Turnock, and B. R. Webber. *Phys. Lett.* **B272**(1991) 368.
11. DELPHI Coll., P. Abreu et al. *Z. Phys.* **C73**(1996) 11.
12. DELPHI Coll., P. Abreu et al. *Z. Phys.* **C54**(1992) 21.
13. DELPHI Coll., P. Abreu et al. *Z. Phys.* **C59**(1993) 21.
14. Y. L. Dokshitzer and B. R. Webber. Cavendish-HEP-97/2, hep-ph/9704298, 1997.